

The origin of the relationship between black hole mass and host galaxy bulge luminosity

C. Martin Gaskell

Astronomy Department, University of Texas, Austin, TX 78712-0259

Abstract. There is a strong decrease in scatter in the $M_{\bullet} - L_{bulge}$ relationship with increasing luminosity and very little scatter for the most luminous galaxies. It is shown that this is a natural consequence of the substantial initial dispersion in the ratio of black hole mass to total stellar mass and of subsequent galaxy growth through hierarchical mergers. “Fine-tuning” through feedback between black hole growth and bulge growth is neither necessary nor desirable.

Keywords: black hole growth, galaxies: fundamental parameters, galaxies: nuclei, galaxies: bulges
PACS: 97.60.Lf, 98.52.Eh, 98.54.-h, 98.62.-g, 98.62.Js, 98.62.Mw, 98.62.Nx, 98.62.Ve, 98.65.Fz

The dispersions in the relationships between black hole mass, M_{\bullet} , host galaxy bulge luminosity, L_{bulge} , and stellar velocity dispersion have been shown [1, 2] to decrease strongly with increasing L_{bulge} (see Fig. 1). The trend in Fig. 1 can easily be modeled by assuming that bulges grow through mergers and that the M_{\bullet}/L_{stars} ratio initially has a log-normal distribution with a substantial dispersion as is observed for the lowest luminosity galaxies. A substantial scatter in the initial M_{\bullet}/L_{stars} ratio is required. “Fine-tuning” through feedback is unnecessary and produces too low a dispersion in M_{\bullet}/L_{bulge} .




FIGURE 1. The $1-\sigma$ scatter in the AGN $M_{\bullet} - L_{bulge}$ relationship as a function of bulge luminosity. Observed points from [2]. The solid line shows the predicted decrease in scatter for hierarchical merging.

REFERENCES

1. Gaskell, C. M. 2009, *Ap.J.* submitted [arXiv:0908.0328].
2. Gaskell, C. M. 2010, in *Co-Evolution of Central Black Holes and Galaxies*, IAU Symposium 267, in press [arXiv:1003.0036].